



Office of Environmental Health & Safety  
333 S. Beaudry Avenue, 28<sup>th</sup> Floor  
Los Angeles, CA 90017  
Phone: (213) 241-3199  
Fax: (213) 241-6816



# SAFETY ALERT

---

No. 14-04

MEDITERRANEAN & ORIENTAL FRUIT  
FLY ERADICATION PROJECT

MARCH 2014  
REV. 1/19/18

---

Between November 8, 2017, and January 3, 2018, one female and two male Oriental Fruit Flies (OFFs), *Bactrocera dorsalis* (Hendel) were trapped in the cities of Rancho Palos Verdes, Lomita, and the community of San Pedro by the California Department of Food and Agriculture (CDFA). Based on the survey data and pest biology, they have determined that there is an Oriental Fruit Fly infestation in the area. They are taking immediate action to eradicate this exotic pest which is known to attack over 250 types of fruits and vegetables.

Over the next several months, the CDFA will implement their action plan to eradicate the fruit fly by employing biological, physical and chemical controls. They will be releasing sterile male fruit flies to prevent the production of offspring. All fruit that shows evidence as a breeding site will be removed and any property within 100 meters of a detection site may be impacted as well. Ground applications of an organic pesticide STATIC Spinosad ME may be applied to properties within a 200-meter radius of where the flies were found.

**At this time, there are no plans to apply any pesticides to District sites.** If additional fruit flies are identified and the CDFA determines a pesticide application is warranted on District property, they will adhere to the District's Integrated Pest Management policy. The project will first be approved and coordinated through the Office of Environmental Health and Safety and Maintenance and Operations (M&O) - Pest Management Unit. In accordance with our Integrated Pest Management policy, a mandatory 72-hour notice will be sent out to all students and staff and a notice will be posted at the impacted site(s).

For additional information, please see the attached fact sheet and area map provided by the CDFA. As they continue to address this issue, additional areas may be added to the treatment zones. Please contact CDFA directly for specific questions regarding this program at 1-800-491-1899. You may also view an information video at the link below:

[http://cdfa.ca.gov/plant/PDEP/treatment/medfly\\_treatment.html](http://cdfa.ca.gov/plant/PDEP/treatment/medfly_treatment.html)

For more general questions or more information on the District's IPM policy please contact OEHS at 213-241-3199 or Facilities, M&O, IPM Unit at 213-745-1435.

### **SUMMARY/BACKGROUND**

- The Oriental fruit fly (OFF) is an invasive pest that attacks over 230 crops: including citrus and other fruits, nuts, vegetables and berries. The short life cycle of the OFF allows rapid development of serious outbreaks, which can cause severe economic losses. Heavy infestations can cause complete losses of crops.
- A great number of crops in California are threatened by the introduction of this pest: including apple, apricot, avocado, bell pepper, fig, grape, grapefruit, lemon, lime, melons, nectarine, orange, peach, pear, persimmon, plum, pomegranate, tangerine, tomato and walnut. In California, the combined 2005 gross value of the commercial hosts potentially affected was over \$9.8 billion.
- Establishment of these flies would cause direct economic losses via damaged fruit, increased pesticide use statewide by commercial and residential growers in efforts to lessen this damage, loss of revenue due to export restrictions on fruit both domestically and internationally, and adverse impacts on native plants through the destruction of their fruit.
- OFF is widespread through much of the mainland of southern Asia and on neighboring islands.
- OFF was first recorded in Hawaii in 1946, and it now attacks nearly every commercial fruit crop grown there except pineapples.
- OFF was first found in California in 1960 and has been reintroduced every year since 1966 through the movement of infested fruits and vegetables into the state. Numerous eradication efforts undertaken by CDFA and USDA over the years have prevented these introductions from becoming permanently established.

### **LIFE CYCLE**

- Female OFF lay eggs in groups of 3 to 30 under the skin of host fruits. A single female can lay more than 1,000 eggs in her lifetime. Maggots tunnel through the fruit, feeding on the pulp. They shed their skins twice and emerge through exit holes in approximately 10 days. The larvae drop from the fruit and burrow two to three cm into the soil to pupate. In 10 to 12 days, the adults emerge. The newly emerged adult females need 8 to 12 days to mature sexually prior to egg laying. With an advanced infestation, breeding is continuous, with several annual generations. Adults live 90 days on the average and feed on honeydew, decaying fruit, plant nectar, bird dung and other substances. The adult is a strong flyer, recorded to travel 30 miles in search of food and sites to lay eggs. This ability allows the fly to infest new areas very quickly.

### **ERADICATION PROGRAM**


- Protecting California's environment from invasive species is the goal, and eradication via male attractant technique is the strategy.
  - Male attractant technique (MAT) is the standard treatment technique for OFF. MAT is conducted in a 1.5-mile radius from each fly find site for a minimum of 9-square-miles.
  - Approximately 600 small, gel-like "bait stations" per square mile are applied to utility poles and street trees at a height of six to eight feet. The technique is repeated every two weeks for a minimum of four applications, or one to two life cycles, depending on the severity of the infestation.
  - These bait stations contain a male attractant (methyl eugenol) that is mixed with a small amount of the pesticide Dibrom (Naled). The bait stations will attract the male fruit flies looking for an opportunity to breed. The females go unmated and no offspring are produced, effectively causing eradication of the population.
  - MAT is specific for this group of flies and will not harm other insects such as bees or butterflies.
  - If larvae, mated females, or numerous males are detected, additional ground treatments may be required to mitigate the spread of OFF. In this case, host trees and plants within a 200-meter radius of the find site are treated with a handheld hose that consists of an organic formulation of Spinosad and protein bait.

### **MORE INFORMATION IS AVAILABLE**

CDFA - [www.cdfa.ca.gov/phpps](http://www.cdfa.ca.gov/phpps)  
CDFA Pest Hotline: 800-491-1899  
APHIS/USDA - [www.aphis.usda.gov](http://www.aphis.usda.gov)



# Oriental Fruit Fly Eradication Project Rancho Palos Verdes, Los Angeles County 2018



- Maximum Program Boundary
- Proposed 200 Meter Treatment Area
- Sensitive Environmental Area/Treatment Mitigations In Place

